

# The effects of convection on the extratropical lower stratosphere

A. E. Dessler


Earth System Science Interdisciplinary Center  
University of Maryland

W. G. Read and J. H. Jiang


Jet Propulsion Laboratory



Plot courtesy of Mark Schoeberl



Plot courtesy of Mark Schoeberl



Evidence for convective transport to 380 K and above:

Fromm et al., Observations of boreal forest fire smoke in the stratosphere by POAM III, SAGE II, and lidar in 1998, *Geophys. Res. Lett.*, 27, 1407-1410, 2000.


Jost et al., In-situ observations of mid-latitude forest fire plumes deep in the stratosphere, *Geophys. Res. Lett.*, 31, L11101, DOI: 10.1029/2003GL019253, 2004.



Ray, E.A., et al., Evidence of the effect of summertime midlatitude convection on the subtropical lower stratosphere from CRYSTALFACE tracer measurements, *J. Geophys. Res.*, 109, D18304, DOI: 10.1029/2004JD004655, 2004.


Livesey et al., Enhancements in lower stratospheric CH<sub>3</sub>CN observed by the Upper Atmosphere Research Satellite Microwave Limb Sounder following boreal forest fires, *J. Geophys. Res.*, 109, D06308, DOI: 10.1029/2003JD004055, 2004.

Wang, P.K., Moisture plumes above thunderstorm anvils and their contributions to cross-tropopause transport of water vapor in midlatitudes, *J. Geophys. Res.*, 108, 4194, DOI: 10.1029/2002JD002581, 2003.


Plot courtesy of Mark Schoeberl







Aura MLS H<sub>2</sub>O VMR  
Aug. 13-Sept. 12, 2004  
380-K pot'l. temp.



MODIS 11- $\mu$ m BT < 210 K  
Aqua + Terra



$$\frac{\partial[X]}{\partial t} + \mathbf{V}_H \cdot \nabla[X] + w \frac{\partial[X]}{\partial z} + \text{convection} = P - L$$



$$\frac{\partial[X]}{\partial t} + \mathbf{V}_H \cdot \nabla[X] + w \frac{\partial[X]}{\partial z} + \text{convection} = P - L$$

Convective  
tendency:

$$\left. \frac{\partial[X]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([X] - [X]_c)$$




Convective  
tendency:

$$\left. \frac{\partial [X]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([X] - [X]_c)$$



$$\left. \frac{\partial [H_2O]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([H_2O] - [H_2O]_{\text{sat}})$$



$$\left. \frac{\partial [\text{H}_2\text{O}]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([\text{H}_2\text{O}] - [\text{H}_2\text{O}]_{\text{sat}})$$




$$\left. \frac{\partial [\text{H}_2\text{O}]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([\text{H}_2\text{O}] - [\text{H}_2\text{O}]_{\text{sat}})$$




$$\left. \frac{\partial [\text{H}_2\text{O}]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([\text{H}_2\text{O}] - [\text{H}_2\text{O}]_{\text{sat}})$$


Convection,  $1/\tau$



$[\text{H}_2\text{O}]_{\text{sat}} - [\text{H}_2\text{O}]$



X



# Single-level isentropic model

- August 13-September 12, 2004; 380 K

$$\frac{\partial[X]}{\partial t} + \mathbf{V}_H \cdot \nabla[X] + \frac{1}{\tau_{BD}}([X] - [X]_{BD}) + \frac{1}{\tau_C}([X] - [X]_C) = 0$$

- Simulated constituent: H<sub>2</sub>O
- Horizontal advection via UKMO winds



# Single-level isentropic model

- August 13-September 12, 2004; 380 K

$$\frac{\partial[X]}{\partial t} + \mathbf{V}_H \cdot \nabla[X] + \frac{1}{\tau_{BD}}([X] - [X]_{BD}) + \frac{1}{\tau_C}([X] - [X]_C) = 0$$

- Simulated constituent: H<sub>2</sub>O
- Horizontal advection via UKMO winds
- Brewer-Dobson circulation parameterized as relaxation



# Single-level isentropic model


- August 13-September 12, 2004; 380 K

$$\frac{\partial[X]}{\partial t} + \mathbf{V}_H \cdot \nabla[X] + \frac{1}{\tau_{BD}}([X] - [X]_{BD}) + \frac{1}{\tau_C}([X] - [X]_C) = 0$$

- Simulated constituent: H<sub>2</sub>O
- Horizontal advection via UKMO winds
- Brewer-Dobson circulation parameterized as relaxation
- Convection parameterized as relaxation



## Fraction of MODIS obs. with 11- $\mu$ m BT < 210 K



$$\frac{\partial [X]}{\partial t} = -\frac{1}{\tau_c} ([X] - [X]_c)$$




# Single-level isentropic model

- August 13-September 12, 2004; 380 K

$$\frac{\partial[X]}{\partial t} + \mathbf{V}_H \cdot \nabla[X] + \frac{1}{\tau_{BD}}([X] - [X]_{BD}) + \frac{1}{\tau_c}([X] - [X]_c) = 0$$


- Simulated constituent: H<sub>2</sub>O
- Horizontal advection via UKMO winds
- Brewer-Dobson circulation parameterized as relaxation
- Convection parameterized as relaxation
  - $1/\tau_\chi$  is proportional to MODIS BT fraction
  - $[X]_c$  is equal to saturation vmr for H<sub>2</sub>O





Model simulation of H<sub>2</sub>O  
VMR at 380 K, Aug. 13-  
Sept. 12, 2004 period






Model simulation of H<sub>2</sub>O  
VMR at 380 K, Aug. 13-  
Sept. 12, 2004 period

Contours are Aura MLS  
H<sub>2</sub>O for that same period





- ✓ Reproduces the mismatch in location of convection and high H<sub>2</sub>O VMRs.




## What about other species?

Convective tendency: 
$$\left. \frac{\partial [X]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([X] - [X]_c)$$

“convective contrast”



Convective tendency:  $\left. \frac{\partial [X]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([X] - [X]_c)$



“convective contrast”

Helium: 5 ppmv, well mixed  
small convective contrast

$$\partial[\text{He}]/\partial t \approx 0$$



Convective tendency:  $\left. \frac{\partial [X]}{\partial t} \right|_{\text{conv}} = -\frac{1}{\tau_c} ([X] - [X]_c)$

“convective contrast”

Species dependent

Helium: 5 ppmv, well mixed  
small convective contrast  
 $\partial[\text{He}]/\partial t \approx 0$



# Conclusions

- Convection can affect the water abundance of the extratropical lower stratosphere to altitudes of 15 km, 380 K
  - Tail of the convective distribution
  - CRYSTAL-FACE data shows the effect drops off rapidly above 380 K
- Different species affected differently
  - “convective contrast” determines effects
  - Important for H<sub>2</sub>O, less so for O<sub>3</sub>
  - Does not mean mass transport is significant
- Dessler, A.E., and S.C. Sherwood (2004), The effect of convection on the summertime extratropical lower stratosphere, *J. Geophys. Res.*, **109**, D23301, DOI: 10.1029/2004JD005209.
  - on the web: <http://www.atmos.umd.edu/~dessler/publications.html>
  - or see me for reprint